

Inflammation Inducible Hybrid Promoters, Vectors Comprising Them And Their Uses

5 The present invention relates to the field of biology. It relates in particular to
the field of the regulation of the expression of genes and, more particularly, it
describes the design and development of a new system for the pharmacological
regulation of the expression of transgenes. The invention is based in particular on
the use of constructs which make it possible to activate the transcription of the
10 transgene in cells, tissues or organs under inflammatory conditions. The invention
thus describes new compositions, constructs and methods allowing the effective
regulation of the expression of a nucleic acid in vitro, ex vivo or in vivo, for example
in chondrocytic cells. The applications which result from the present invention are
many, in the experimental, clinical, therapeutic or diagnostic fields, for example.

15 Controlling the level and the duration of the expression of the transgenes is
necessary for many applications. Thus, gene therapy, the production of recombinant
proteins in vitro, the construction of transgenic animals, the study of the effects of a
gene or of the bioavailability of a protein and the like are as many situations in which
an appropriate control of genetic expression can be implemented and can bring
20 improvements.

 Gene therapy is aimed more particularly at expressing genes (or nucleic
acids) which are either defective in a given individual, or which are of therapeutic
interest in the context of a specific pathology. The expression of these genes
25 requires, apart from the vectorization procedure, an induction with a promoter which
binds RNA polymerase and the transcription factors required to obtain the synthesis
of messenger RNA and therefore protein synthesis. Noninducible viral or eucaryotic

promoters are frequently used in gene therapy to express a gene at the level of a tissue. Although it is easy to carry out, this approach does not allow the regulation of the expression of the transgene according to the environment and the external stimuli. Furthermore, the gene directed by a viral promoter will be expressed equally well in all the cells into which the vector would have penetrated, without any targeting. Finally, experimentalists have sometimes been confronted with the weakness of the expression of the viral promoter *in vivo* in relation to that observed *in vitro*.

Various artificial regulators of transcription have been designed in the prior art, which are activated by a xenobiotic molecule which binds to the promoter sequences for transcription of the transgene.

A first illustration of these regulators was constructed by fusion of the *E. coli* Lac repressor with the herpes simplex virus (HSV) VP16 transactivator domain. Two versions of these regulators exist, one capable of being activated by isopropyl β -D-thiogalactoside (IPTG) and the other inactivated by IPTG (Bain S. et al., *Proc Natl Acad Sci USA*, **88** (1991) 5072-5076; Labow M. et al., *Mol. Cell. Biol.*, **10** (1990) 3343-3356).

Another system was constructed by fusion of the *E. coli* Tet repressor with the HSV VP16 transactivator domain. Two versions of these regulators also exist, one capable of being activated by tetracycline or its derivatives and the other inactivated by these same molecules (Gossen M. and Bujard H., *Proc Natl Acad Sci USA*, **89** (1992) 5547-5551; Gossen M. et al., *Science*, **286** (1995) 1766-1769).

Another system was constructed by fusion of the DNA-binding domain of the *S. cerevisiae* GAL4 protein with the ligand-binding domain of the human progesterone receptor and the HSV VP16 transactivator domain; this version is activated by a progesterone analogue such as RU486 (Wang Y. et al., *Proc Natl*

Acad Sci USA, **91** (1994) 8180-8184). A fusion of the drosophila ecdysone receptor with the HSV VP16 transactivator domain has also been described, activated by ecdysone and the analogues of this steroid hormone (No D. et al., *Proc Natl Acad Sci USA*, **93** (1996) 3346-3351). Another system takes advantage of the capacity of certain immunosuppressive molecules (cyclosporin A, rapamycin and its derivatives) to promote the combination of certain cellular proteins. A transcriptional regulator then consists of two protein subunits; the first may be formed by the fusion of a chimeric DNA-binding domain and of three copies of the human FKBP protein and the second by the fusion of the rapamycin-binding domain of the human FRAP protein and of the transactivator domain of the human NFkB p65 subunit. This transcriptional regulator is activated by rapamycin which allows the dimerization of the two subunits (Rivera V. et al., *Nat. Med.*, **2** (1996) 1028-1032).

Although these systems make it possible to obtain satisfactory levels of regulation in some tissues, they exhibit, nevertheless, certain disadvantages which limit their conditions of use. Thus, these transcriptional regulators are xenogenic proteins in humans. They indeed consist of protein fragments obtained from bacteria, viruses, yeasts or insects or, when the protein domains are of human origin, their joining creates sequences which are foreign to humans. These protein domains may therefore induce a cytotoxic immune reaction, causing the destruction of the cells which express the gene of interest under the control of the xenogenic transcriptional regulator, and thus the termination of the expression of the transgene. This situation may necessitate the use of repeated administrations of the therapeutic gene, which constitutes a major disadvantage, in particular when this involves a traumatic surgical act, and which is not always effective, in particular when the vector of the therapeutic gene is a virus whose first injection causes an immune reaction. In addition, the levels of expression observed with the prior art systems of regulation are not always satisfactory.

The present invention now describes a new improved system for the controlled expression of nucleic acids in cells. This invention provides a hybrid promoter which is inducible by the principal mediators of inflammation and by exogenous compounds. This promoter preferably leads to the expression of a gene at the level of the tissue which is the seat of an inflammation, in response to the mediators synthesized during the inflammatory process. The principal benefit of this promoter lies in the possibility of an expression of the gene-medicament which can be modulated according to the intensity of the inflammation. Furthermore, the expression of the gene maybe increased if NSAIDs are being taken. This is therefore an approach which is transposable to many inflammatory situations for which a gene therapy is envisaged.

The present invention consists more particularly in the construction of a hybrid promoter comprising a module for responding to the PPARs and a module for responding to the cytokines produced during inflammation. This second module comprises more particularly sequences derived from the promoter of the gene for type IIA-1 nonpancreatic secreted phospholipase A2 (PLA2s), which is expressed in some human cells, in particular the chondrocytes, under inflammatory conditions. The combination of these two modules makes it possible to generate a promoter allowing a specific expression of genes under inflammatory conditions, which expression may be increased by exogenous anti-inflammatory medicaments or mediators. As will be illustrated in the examples, this hybrid promoter preferably possesses a low basal activity, but a very high activity under induction conditions. This type of promoter is therefore particularly suitable for the expression of products having an anti-inflammatory activity, as will be explained in detail in the remainder of the text. The promoters of the invention are most particularly advantageous for the expression of genes in articular tissues, in particular the chondrocytes, in different

pathological situations in which an inflammatory component is present, and in particular osteoarthritis.

Osteoarthritis is a disease of aging which is accompanied by articular destruction which causes a variable degree of invalidity (Chevalier X, 1998). The treatment of this pathology is currently considered as a priority in terms of public health because of its frequency and the handicap which it causes. From a physiopathological point of view, even though the mechanisms of initiation still remain poorly understood, the tissue target responsible for the articular destruction is the cartilage, a tissue composed of an extracellular matrix rich in type II collagen and in proteoglycans and articular chondrocytes, sole cartilage cell type, which are responsible for the matrix synthesis. The chondrocytes also secrete growth factors and natural protease inhibitors (anabolic activity) as well as proteases (catabolic activity). Physiologically, there is an equilibrium between these activities (cartilaginous homeostasis). During osteoarthritis, there is a disequilibrium at the expense of the anabolic activity because of the stimulation of the synthesis of proteases (essentially metalloproteases) and an inhibition of the synthesis of inhibitors of metalloproteases (TIMP or tissue inhibitor of metalloproteases) (Lefebvre V et al., 1997; Dean J et al., 1990; Hornebeck W, 1990). These modifications in synthesis are due to the presence of pro-inflammatory mediators produced by the chondrocytes and found in large quantities in the synovial fluids of patients suffering from osteoarthritis or from rheumatoid arthritis, cytokines (IL-1, TNF) and lipid mediators, in particular prostaglandin E2. This prostaglandin stimulates the synthesis of collagen in low concentration like the growth factor but inhibits in high concentration (Di Battista JA et al., 1996).

The present invention now provides hybrid transcriptional promoters which are particularly suitable for the expression of therapeutic genes in articular pathologies

such as osteoarthritis. In addition, the invention also describes constructs allowing screening of the transfer and/or of the expression of a gene of interest into the chondrocytes, further improving the advantageous properties of the constructs for this type of applications.

A first subject of the invention consists more particularly in a hybrid promoter comprising :

- a) a PPAR response element and
- b) the whole or part of the promoter of the PLA2s IIA gene.

The present invention consists in particular in the demonstration that this type of construct produces a synergistic effect on the expression activity, without impairing the regulated and specific (or preferential) nature of inflammatory situations. The invention also describes vectors, plasmids and compositions comprising these promoters, as well as their uses for the expression of genes in vitro, ex vivo or in vivo.

The first element of the hybrid promoters of the invention therefore comprises a PPAR response element. PPARs (Peroxisome Proliferator-Activated Receptors) belong to the superfamily of nuclear receptors comprising the receptors for steroid hormones, the receptors for thyroid hormones, for retinoic acid and for vitamin D (Schoonjans K et al., 1996). PPARs exist in three isoforms: PPAR α , PPAR β and PPAR γ . The PPARs bind to specific sequences at the level of the promoters of target genes (PPRE sequences), and induce the expression of these genes.

A PPAR response element (PPRE, "Peroxisome Proliferator Response Element") is therefore a nucleic acid region capable of binding a PPAR, it being

possible for the binding of the PPAR to then mediate a signal around neighboring nucleic regions, in particular the regions derived from PLA2s according to the invention. A PPAR response element according to the invention is more particularly a nucleic acid region capable of binding a PPAR, whether it is a PPAR α , β or γ . It is still more preferably a nucleic acid capable of binding a PPAR α or a PPAR γ . For carrying out the invention, the PPAR response element comprises more particularly one or more PPAR-binding sites. Such binding sites have been described in the prior art, such as, for example, in different human promoters (consensus site DR1, gene for apolipoprotein AII, for example). Such sites may also be artificially constructed, and tested for their PPRE properties, as is described below.

In a specific embodiment of the invention, the PPAR response element comprises one or more sites having the sequence CAAAACTAGGTCAAAGGTCA (SEQ ID NO:1) or functional variants of this sequence. The sequence SEQ ID NO: 1 corresponds to the consensus region DR1.

The term functional variant designates any modified sequence conserving the properties of PPRE as mentioned above, that is to say in particular the capacity to bind a PPAR. The modifications may comprise one or more additions, mutations, deletions and/or substitutions of nucleotides in the sequence considered. These modifications may be introduced by conventional molecular biology methods, such as in particular site-directed mutagenesis or, more practically, by artificial synthesis of the sequence in a synthesizer. Generally, the variants conserve at least 50% of the residues of the initial sequence indicated. More preferably, the variants possess modifications affecting less than 8 nucleotides in the sequence considered. The variants thus obtained are then tested for their PPRE activity. This property may be checked in various ways, and in particular:

(i) by bringing the test sequence into contact with a PPAR (for example in an acellular test), and detecting the formation of a complex (for example by gel migration retardation);

(ii) by inserting the test sequence into an expression cassette comprising a minimal promoter and a reporter gene, introducing the cassette into a cell, and detecting (where appropriate assaying) the expression of the reporter gene in the presence and in the absence of a PPAR and of a ligand for a PPAR;

(iii) by any other technique known to a person skilled in the art, making it possible to detect the interaction between a nucleic acid and a protein, for example.

A variant is considered to be functional for the purpose of the present invention when the activity measured, for example in (ii) above, is preferably at least equal to 50% of that measured with a site having the sequence SEQ ID NO:1, more preferably at least equal to 75%. Functional variants of PPAR-binding sites for the purpose of the invention are described, for example, in Juge-Aubry et al. (J. Biol. Chem. 272 (1997) 25252) and in Nakshatri et al. (NAR 26 (1998) 2491), which are incorporated into the present application by way of reference.

Specific examples of PPAR response elements according to the invention are represented by the elements (DR1)₂ 17 (SEQ ID NO :2), (DR1)₂ 21 (SEQ ID NO :3) and (DR1)₂ 31 (SEQ ID NO :4) as described in the examples, or functional variants thereof.

Another variant is a fragment having the sequence SEQ ID NO :1, comprising residues 8-20 thereof.

The sequences SEQ ID NO :1-4 according to the invention represent PPREs essentially corresponding to PPAR α and γ . As illustrated in the examples, these sequences possess high inducibility by the PPARs, linked in particular to the particular arrangement of the binding sites. In this regard, a particular subject of the

invention also consists in a nucleic acid comprising a sequence chosen from SEQ ID NOS :2-4.

According to another particular embodiment, the PPAR response element
5 comprises one or more J sites of the human apoAII promoter (nucleotides -734 to -716) or functional variants of this sequence, which corresponds to activation by the PPAR β s.

As indicated above, in the compositions according to the invention, the PPAR
response element may comprise several sites for binding to a PPAR. This may be a
10 repetition of the same site, or combinations of different sites, the repetition of identical sites being preferred. More particularly, the response element comprises up to 30 binding sites, preferably from 3 to 20, more preferably from 1 to 5. A preferred embodiment of the invention is a construct comprising 2 DR1 binding sites, the results presented in the examples showing the advantageous properties of such
15 constructs in terms of induction and of levels of expression, in particular in the chondrocyte cells (SEQ ID NO : 2-4).

For the preparation of a hybrid promoter according to the invention, the PPAR
response element (a) is combined with the whole or part of the PLA₂s IIA-1 promoter
20 (element b)).

The nonpancreatic secreted phospholipase A₂ is a protein activated by
interleukin-1 beta, and leads to the production of inflammation mediators, in
particular lipid mediators (prostaglandin E₂). In particular, prostaglandin PGE₂ is
derived from the activation, by interleukin-1 (IL-1), of nonpancreatic secreted
25 phospholipase A₂ (PLA₂s) (Nevalainen TJ et al., 1993; Hulkower KJ, 1997) and releases arachidonic acid from phospholipids and of inducible cyclooxygenase (COX-2) (Amin AR, 1997; Geng Y et al., 1995; Knott et al., 1994), the latter converts

the arachidonic acid to PGH₂. PGH₂ is then converted to PGE₂ by PGE synthase. Two types of phospholipase A₂ have been identified in human cells, in particular the chondrocytes. The nonpancreatic secreted phospholipase A₂ (PLA_{2s}) is a 14 kDa protein which is abundantly synthesized and secreted in inflammatory synovial fluids (Seilhamer JJ et al., 1989). Cytosolic phospholipase A₂ (PLA_{2c}) is a cytosolic protein of 85 Kda whose activation occurs via its phosphorylation by the MAP kinases and its translocation to the membrane (Clark JD et al., 1991). IL-1 induces the expression of the COX-2 and PLA_{2s} genes without affecting that of the PLA_{2c} gene in the chondrocytes (Berenbaum F et al., 1994; Jacques C et al., 1997). The levels of secreted PLA₂ in the joint and in the serum are correlated with the level of inflammation, in particular the level of osteoarthritis in the case of the chondrocytes (Pruzanski W et al., 1994; Lin M et al., 1996). The cooperation between PLA_{2s} and COX-2 results in a sustained production (>48 hours) of prostaglandin PGE₂ under IL-1 β stimulation (Bingham CO et al., 1996; Kuwata H et al., 1998) whereas PLA_{2c} and the constitutive cyclooxygenase COX-1 would be responsible for an early response. In our previous studies, a correlation had been shown between the synthesis of PGE₂ and the level of activity of PLA_{2s} linked to the chondrocytic membranes (Jacques C et al., 1997).

The present invention now proposes using the whole or part of the promoter of the PLA_{2s} gene for regulating the expression of nucleic acids of interest, in therapeutic applications in particular. The invention proposes in particular using the whole or functional parts of the PLA_{2s} IIA promoter for the construction of a regulated hybrid promoter active in cells exhibiting an inflammatory situation. In this regard, the applicants have shown that a fragment of 247 base pairs upstream of the first exon of the human gene for PLA_{2s} IIA (SEQ ID NO :5) was responsible for a stimulation 6 to 8 times the activity of the CAT reporter gene by IL1 β in rabbit chondrocytes in primary culture. Three regulatory regions were characterized in this

promoter: the proximal region (-85/-114) ensures a basal transcriptional activity and binds the factor Sp1; the distal region (-247/-210) binds members of the NF1 family and the receptor for the glucocorticoids, which stabilize the binding of the C/EBP factors between positions -198/-190 relative to the site of initiation of transcription.

- 5 The family of C/EBP factors comprises three principal activating members C/EBP α , C/EBP β and C/EBP δ . The factors C/EBP β and C/EBP δ are activated at the transcriptional and post-transcriptional level by the pro-inflammatory cytokines. They are expressed in the chondrocytes whereas C/EBP α is not. C/EBP β and C/EBP δ are responsible for the activation of the transcription of the gene for sPLA2-IIA by IL1 β .
- 10 The applicants have now shown that they are also critically involved in the induction of the transcription of the gene for COX-2 by IL-1 β in human and rabbit chondrocytes.

In a specific embodiment, element b) of the hybrid promoters of the invention comprises the whole or part of the sequence SEQ ID N :5 or of a functional derivative thereof. The sequence SEQ ID NO :5 corresponds to nucleotides -247 to +20 of the human gene for PLA2s IIA (residues 5 to 271).

15

The term « part » designates more particularly any fragment of the sequence considered which conserves biological functionality. This is more particularly a part ensuring a basal transcriptional activity for the hybrid promoter (TATA box, for example) and a character regulatable by the inflammation mediators, in particular by interleukin-1 β .

20

In this regard, a preferred variant of the invention relates to a hybrid promoter comprising at least part of the PLA2s IIA promoter conferring induction by interleukin-1 β .

25 More particularly, element (b) of a hybrid promoter according to the invention may comprise the following sequences:

- at least residues 51 to 61 of SEQ ID NO :5 (corresponding approximately to residues -200/-191 of the human PLA2s IIA gene), involved in the regulation by IL1, and/or,

- at least residues 23 to 32 of SEQ ID NO :5 (corresponding approximately to residues -229/-220 of the human PLA2s IIA gene), representing a half site NF1 and a half site GRE (these sequences are involved in the regulation by certain compounds such as dexamethasone), and/or

- at least residues 148 to 155 of SEQ ID NO :5 (corresponding approximately to residues -104/-97 of the human PLA2s IIA gene), involved in the binding of Sp1, and/or

- at least residues 5 to 170 of SEQ ID NO :5 (corresponding approximately to residues -247/-85 of the human PLA2s IIA gene), or alternatively

- at least residues 51 to 170 of SEQ ID NO :5 (corresponding approximately to residues -200/-85 of the human PLA2s IIA gene).

A specific example of a hybrid promoter comprises a PPAR response element (DR1)₂ 21 linked to fragment -247/+20 of the PLA2s IIA gene. Such a promoter comprises the sequence SEQ ID NO :6, which is also a specific subject of the present invention. In this sequence, the PPRE element essentially corresponds to residues 13-53 and the PLA2s element to residues 62-332. Residues 1-12 and 54-61 correspond to neutral regions from the functional point of view, resulting from clone steps and making it possible to modify the structure of the promoter (cloning sites).

Of course, sequence variations may occur in the regions considered, without substantially affecting the transcriptional activity of the hybrid promoter of the invention (mutations, deletions, substitutions, insertions and the like), in particular on

one to five bases. It is understood that such variants represent specific embodiments of the present invention.

The PPAR response element (PPRE) and the region derived from PLA2s IIA are functionally arranged in the hybrid promoter of the invention, that is to say such that the hybrid promoter exerts a transcriptional activity in the presence of inflammation mediators, preferably regulated by the PPRE element.

Preferably, element a) is positioned upstream (in 5') of element (b) in the hybrid promoter (cf. Figure 1). This configuration indeed makes it possible to ensure a high and regulated level of expression in the cells, in the presence of inflammation mediators (LTB4) and/or of PPAR activators.

The different functional domains above can be linked directly to each other, or separated by or combined with nucleotides which do not significantly affect the regulated nature of the promoter. Such nucleotides may be residues which are neutral from the functional point of view, resulting for example from cloning steps (PCR ends, restriction sites and the like). These nucleotides may also possess biological properties, making it possible to confer improved characteristics or performance on the system of the invention (enhancer of housekeeping genes, tissue-specific enhancer, silencer, intron, splicing site and the like). In this regard, in a particular embodiment of the invention, the hybrid promoter comprises, in addition, an element c) conferring tissue specificity. This element c) may be located between elements a) and b), or downstream (3') of element b). The term « tissue specificity » designates a preferential or predominant expression in a given (type) of cells, without however completely excluding a residual or lower expression in other cells.

A specific embodiment of the invention consists more specifically in a hybrid transcriptional promoter comprising, in the 5'->3' orientation :

- a) a PPAR α and/or γ response element preferably comprising the whole or part of a sequence SEQ ID NO :1 to 4,
- b) the whole or part of the sequence SEQ ID NO :5, and
- c) where appropriate, an element conferring tissue specificity.

5

In a specific application, element c) confers specificity of expression for the chondrocytic cells, and in particular it comprises the whole or part of the sequence SEQ ID NO :7 or of a variant thereof. This type of promoter is particularly suitable for the regulated and selective expression of nucleic acids in the chondrocytes, in particular in an inflammatory situation (osteoarthritis).

10

The cartilaginous matrix consists of a large number of collagenous and noncollagenous proteins. Some genes encoding proteins of this matrix are specifically expressed by the chondrocyte like the genes for type II collagen, aggrecan, matrilin-1 (or cartilage matrix protein) and chondroadherin. The latter two

15 proteins interact specifically with integrins synthesized by the chondrocytes and are responsible for anchoring of the cell to the other components of the matrix (Makihira S et al., 1999; Camper L et al., 1997). The expression of the genes for type II collagen and aggrecan have been widely cited in the literature as positive markers of chondrocytic differentiation whereas the synthesis of type I collagen marks a

20 phenomenon of dedifferentiation (Benya P et al., 1986). The group of Pr de Crombrugghe has identified a sequence of 18 base pairs which is necessary and sufficient to direct the expression of type II collagen in the chondrocytes. This sequence binds a nuclear factor (SOX9) which is moreover thought to be involved in chondrocytic differentiation (Mukhopadhyay K et al., 1995; Zhou G et al., 1995). This

25 factor is colocalized with the expression of type II Collagen in the primary chondrocytes. It binds to the sequence CATTCAT at the level of the promoter of the type II collagen gene and activates its transcription.

For the gene of interest to no longer be expressed only in an inducible manner but also specifically in the chondrocyte, elements of the promoter of type II collagen which are capable of binding the SOX factors (SOX9, SOX6 and L-SOX5) (Mukhopadhyay K et al., 1995; Zhou G et al., 1995; Lefebvre V et al., 1997) may be integrated. A nucleotide sequence described by Zhou et al. (1995) and Lefebvre V et al. (1997) formed of a duplicate of 231 bp of the first intron of type II collagen may be used. This construct proved to be the most effective in the system. It can be inserted downstream of the hybrid construct (DR1)2 21 -promoter of sPLA2 IIA, by including splice donor (SD) and acceptor (SA) sites on either side, this being with the aim of constituting a first synthetic intron following the first exon of 20 untranslated base pairs of the gene for sPLA2-IIA as schematically represented in Figure 7A. The insertion of a first intron generally facilitates the expression of the transgene situated downstream. This insertion would make it possible, on the one hand, to restrict and, on the other hand, to enhance the expression of the transgene in the chondrocytes by virtue of the presence of the site for the Sox9 factor. In another construct, the sequence containing the sites for binding the Sox factors as a promoter may be inserted before the sites for binding the PPARs as schematically represented in Figure 7B.

Another subject of the invention relates to any nucleic acid comprising a hybrid promoter as defined above and a gene of interest. The gene of interest may be any nucleic acid (cDNA, gDNA, synthetic or semisynthetic DNA, RNA, and the like), of diverse origin (of animal, human, plant, bacterial, viral or artificial origin, and the like), encoding a product of interest (RNA, polypeptide or peptide). It is advantageously a nucleic acid encoding an anti-inflammatory product, that is to say a product capable of restoring or of reducing the inflammatory phenomena in a cell, tissue or organ. Among the anti-inflammatory products, there may be mentioned in particular TIMP or

other cytokines or enzymes capable of reducing the inflammatory phenomenon. The product of interest may be, more generally, any enzyme, hormone, growth factor, antibody, immunogenic peptide, lipoprotein, toxin, antibody or antibody fragment, antisense, transcription factor and the like, of therapeutic or vaccinal interest.

5 The gene of interest is generally placed downstream (3') of the hybrid promoter, under the transcriptional control of the latter. Thus, a preferred nucleic acid for the purposes of the invention comprises, in the 5'->3' orientation :

a) a PPAR α and/or γ response element preferably comprising the whole or part of a sequence SEQ ID NO :1 to 4,

10 b) the whole or part of the sequence SEQ ID NO :5,

c) where appropriate, an element conferring tissue specificity, and

d) a gene encoding a product of interest, in particular of therapeutic or vaccinal interest.

15 Another subject of the invention consists in a vector comprising a hybrid promoter or a nucleic acid as defined above. The vector may be of varied, in particular plasmid, episomal, chromosomal, viral or phage, nature and/or origin, and the like. Preferably, the vector is a plasmid or a recombinant virus, more preferably a plasmid.

20 As an example of a viral vector, there may be mentioned in particular a recombinant adenovirus, a recombinant retrovirus, an AAV, a herpesvirus, a vaccinia virus and the like, whose preparation may be carried out according to methods known to persons skilled in the art.

As regards the plasmids, there may be mentioned any replicative or
25 integrative plasmid which is compatible with use in mammalian, in particular human, cells.

Another subject of the invention consists in a composition comprising a nucleic acid or a vector as defined above. The composition may contain, moreover, any vehicle acceptable for administration in vivo or ex vivo of constructs (in particular any pharmaceutically acceptable vehicle), and/or for a stable storage thereof. As vehicle, there may be mentioned in particular isotonic, buffered, saline solutions optionally comprising additives such as high-molecular weight proteins or polymers. Moreover, the compositions of the invention may comprise agents which facilitate the penetration of the genetic constructs of the invention into cells. There may be mentioned in particular lipids, polymers, peptides and the like, which make it possible to improve cellular transfection.

To this effect, a particular composition according to the invention comprises one or more polycationic lipids, optionally coupled to polycationic polymers. This type of formulation is particularly suitable for the transfer of genes into the chondrocytes, which are surrounded by a matrix rich in negatively charged proteoglycans. Polycationic lipids can therefore become easily integrated into this matrix. The addition of polycationic polymers such as PEI (polyethylenimine) comprising incompletely protonable amines at the physiological pH makes it possible to increase the efficiency of transfection of the chondrocytes. These amines function as a sponge for protons with endosomolytic properties. The combination consisting of the cationic lipids, the PEI and the plasmid carrying the transgene constitutes a particular composition which is preferred in the context of the present invention.

Moreover, the compositions of the invention may also comprise one or more targeting elements which make it possible to ensure a preferential transfection toward desired cell types. In this context, the targeting elements may consist of ligands for specific receptors, receptors for ligands, antibodies or antibody fragments, and the like.

For targeting toward chondrocytic cells, a preferred composition for the purposes of the invention comprises, in addition, one or more molecules which possess affinity for the chondrocytic membrane. Among these proteins, there may be mentioned in particular the Chondroadherin protein, which interacts with a good affinity with integrin alpha2 beta1 of the chondrocytes (Camper L et al., 1997). It is a small protein of 36 kDa, rich in leucine. This protein may be produced (for example in a recombinant baculovirus), purified and incorporated into a composition (in particular a liposome) of the invention. The hydrophobicity of the protein can, moreover, be increased by addition of a farnesylation sequence at the C-terminal like the Q-Ras4B end (Leevers SJ et al., 1994; Stokoe D et al., 1994; Zlatkine et al., 1997).

It is understood that any other targeting element may be envisaged, depending on the desired applications.

Moreover, another subject of the invention further consists in a composition comprising (i) a nucleic acid or a vector as defined above and (ii) a PPAR activator, for use simultaneously, separately or spaced out over time.

As indicated above, the hybrid promoters comprise a PPAR response element. This PPAR may be released (or induced) into the cells, for example in an inflammatory situation, or also in the presence of exogenous activators (i.e. added or administered). In this context, the expression « for use simultaneously, separately or spaced out over time » indicates that the vector and the activator may be prepared separately, packaged separately and used sequentially, to allow control of the expression of the nucleic acid of interest. Typically, the vector and the activator are packaged separately and used spaced out over time, the combination of these different elements in a cell, tissue or organ, and the like, leading to the desired effect of regulating or enhancing expression.

Depending on the hybrid promoter or the desired applications, various types of ligands, natural or synthetic, may be used. Thus, this may be preferably a PPAR γ activator and/or a PPAR β activator and/or a PPAR α activator.

The PPAR γ -activating ligands may be chosen from natural or synthetic ligands. As natural ligands, there may be mentioned fatty acids and eicosanoids (for example linoleic acid, linolenic acid, 9-HODE, 5-HODE) and, as synthetic ligands, there may be mentioned thiazolidinediones, such as in particular rosiglitazone (BRL49653), pioglitazone or troglitazone (see for example Krey G. et al., *Mol. Endocrinol.*, 11 (1997) 779-791 or Kliewer S. and Willson T., *Curr. Opin. in Gen. Dev.* 8 (1998) 576-581), the compound RG12525 or 15 dioxo- Δ 12-14 PGJ2.

The PPAR α -activating ligands are, for example, the fibrates such as fibric acid and its analogs. As analogs of fibric acid, there may be mentioned in particular gemfibrozyl (*Atherosclerosis* 114(1) (1995) 61), bezafibrate (*Hepatology* 21 (1995) 1025), ciprofibrate (*BCE&M* 9(4) (1995) 825), clofibrate (*Drug Safety* 11 (1994) 301), fenofibrate (*Fenofibrate Monograph*, Oxford Clinical Communications, 1995), clinofibrate (*Kidney International*. 44(6) (1993) 1352), pirinixic acid (Wy-14,643), 5,8,11,14-eicosatetranoic acid (ETYA), or the nonsteroidal anti-inflammatory agents (NSAID) such as ibuprofen and indomethacin. These different compounds are compatible with a biological and/or pharmacological use in vitro or in vivo.

Moreover, the compositions according to the invention may comprise several PPAR activators in combination, and in particular a fibrate or a fibrate analog combined with a retinoid.

These different ligands may be used at conventional doses which are described in the prior art and are illustrated in the examples.

The compositions according to the invention may be formulated in any suitable type of material, such as a tube, ampoule, bag, vial, syringe and the like, and stored in the cold (or frozen) or in freeze-dried form.

The invention may be used for expressing a gene in various types of cell, tissue or organ, in vitro, ex vivo or in vivo. In particular, this may be a mammalian, preferably human, cell, tissue or organ, in particular in an inflammatory state. By way of illustration, there may be mentioned chondrocytic cells (or cells of the bone matrix) muscle cells (or a muscle), hepatic cells (or the liver), cardiac cells (or the heart, the arterial or vascular wall), nerve cells (or the brain, the marrow and the like) or tumor cells (or a tumor).

Preferably, the constructs, compositions and method of the invention are used for the regulated expression of a nucleic acid in a chondrocytic cell in vitro, ex vivo or in vivo. The results presented in the examples illustrate more particularly the advantages of the invention in vivo or in vitro in this type of cells.

The invention also relates to the use of a vector as defined above for the preparation of a composition intended for inducing the expression of a gene in a tissue in an inflammatory situation. It also relates to a method for the expression in a gene in a tissue in an inflammatory situation, comprising the administration of a vector or of a composition above to a subject.

Advantageously, it involves a use or a method for the expression of a gene in a chondrocyte.

The invention also relates to the use of the above nucleic acids, vectors or compositions for the treatment of osteoarthritis.

For a use in vitro or ex vivo, the cells may be brought into contact with the compositions or vectors of the invention according to various protocols. Thus, the cells in culture may be incubated directly with the vectors or compositions of the invention, for example with a vector comprising the hybrid promoter and a gene of

interest, where appropriate in the presence of the activator. Alternatively, the cells may be incubated in a first stage with the vector or nucleic acid and then, in a second stage (after culture and optionally selection of the modified cells), the activator may be added. These experiments may be carried out in any appropriate medium and device, preferably in a plate, dish, flask, under sterile conditions. The quantities of cells, vector and ligand can be easily adapted by persons skilled in the art, on the basis of the information provided in the examples and of their general knowledge.

For a use in vivo, the cells (or organs, tissue and the like) are brought into contact by administration of the nucleic acids, vectors or compositions in vivo, simultaneously, separately or spaced out over time. To this effect, the vectors, compositions or nucleic acids are generally administered by the parenteral, in particular intramuscular, intravenous, subcutaneous, intradermal, intratumoral, intrabone, intraarticular or stereotaxic route. The choice of the mode of administration may be guided by the application envisaged, the tissue targeted and/or the type of product of interest encoded by the transgene. For this administration, the compositions of the invention may comprise any agent promoting cellular transfection (cationic polymer, lipid and the like).

The activator may be administered before, simultaneously or after the vectors or nucleic acids. In this regard, the administration of the ligand may be carried out by the oral, anal, intravenous, intraperitoneal, intraocular or intramuscular route, for example.

The doses used may be adapted by persons skilled in the art, on the basis of the in vivo data published in the literature. Thus for example, for a form not soluble in water, typical doses of ligand such as BRL 49653 are between 5 and 50 mg/kg, for example 30 mg/kg, which make it possible to obtain a plasma concentration close to about 15 µg/ml at least. For a water-soluble form of ligand, whose bioavailability is greater (for example a maleate salt of BRL49653), the typical doses are lower,

generally less than 5 mg/kg, for example from 0.01 to 1 mg/kg. These doses can be quite obviously adapted by persons skilled in the art as a function of the constructs used, the ligands used, and the desired applications and effects. Moreover, repeated administrations of ligand may be carried out.

5 In general, the doses of vector used may vary between 0.01 and 1000 μ g, or more, depending on the desired applications.

10 In a typical experiment, a vector according to the invention is administered by the local route (directly into the affected tissue, for example the seat of inflammation, in particular by the intraarticular route) in the presence of a transfecting agent. Under the effect of the mediators released by the cells (PG2, IL1, LTB4, and the like), an expression of the gene of interest is observed which may be stimulated or enhanced by administration of one or more PPAR ligands, as defined above, or in the presence of endogenous PPARs.

15 Thus, the release of arachidonic acid and of other fatty acids by phospholipases A₂ results in the synthesis of other icosanoids and lipid mediators. These mediators activate the transcription factors PPARs (Peroxisome Proliferator-Activated Receptors). These factors are also activated by certain medicaments such as hypolipemic fibrates, the thiazolidinediones used in diabetes and the nonsteroidal
20 anti-inflammatory drugs (NSAIDs) (Lehmann JM et al., 1997). The chondrocytes express the PPAR α and PPAR γ factors (Bordji et al., J. Biol. Chem., in press). The icosanoids (derivatives of arachidonic acid) such as the prostaglandins (PGE2, PGJ2) induce PPAR γ (Kliwer SA et al., 1995; Krey G et al., 1997); in particular, prostaglandin 15 dioxy Δ 12-14 PGJ2 is the most potent activator of PPAR γ . The
25 leukotriene LTB4 and the fibrates activate PPAR α (Devchand RP et al., 1996) and the long fatty acids activate the two isoforms PPAR α and γ .

The results presented in the examples illustrate the synergistic properties of the hybrid promoters, in the presence of two activating signals.

5 The invention also relates to any cell modified by bringing into contact with a composition or a vector as defined above, especially any mammalian, in particular human, cell, more specifically chondrocytes.

10 The invention also relates to various fragments of the PLA2s promoter as described above, which can be used for example to confer a regulatable character to promoters.

The present invention will be described in greater detail with the aid of the following examples which should be considered as illustrative and nonlimiting.

15 **LEGEND TO THE FIGURES**

Figure 1 : Schematic representation of a hybrid promoter of the invention controlling the expression of TIMP1.

20 Figure 2 : Gel retardation binding analysis. The radiolabeled sequences DR1, (DR1)₂ 17 , (DR1)₂ 21, (DR1)₂ 31 are incubated with 5 or 10 µl of Cos-1 extracts enriched with PPAR_γ and RXR. Lanes 1, 4, 7, 10 correspond to 5 µl and lanes 2, 5, 8, 11 to 10 µl of extracts. Lanes 3, 6, 9 and 12 correspond to 10 µl of nontransfected Cos-1 extracts.

25 Figure 3 : Quantification of the radioactivity obtained in the dimeric and tetrameric bands and the free probe (PPAR_γ).

Figure 4 : Quantification of the radioactivity obtained in the dimeric and tetrameric bands and the free probe (PPAR α).

Figure 5 : Effect of PGJ2 (PPAR γ inducer) and Wy-14643 (PPAR α inducer) on the synthetic promoters.

- 5 Figure 6 : The rabbit chondrocytes were transfected with the plasmid containing the element (DR1)₂ 21- (-247/+20)sPLA2-CAT. The different inducers mentioned above are added to the culture medium.

Figures 7A and 7B : Schematic representation of the constructs into which elements of the type II collagen promoter capable of binding the SOX factors have been
10 integrated respectively downstream and upstream of the hybrid construct (DR1)₂ 21 -promoter of sPLA2 IIA.

MATERIALS AND METHODS

1- Cell cultures and transfections.

- 15 The hind and forelegs of 3-week-old prepubescent rabbits (Fauve de Bourgogne) were dissected and chopped up under sterile conditions. The cartilage is removed with a scalpel and then the chondrocytes are dissociated by means of proteolytic enzymes: hyaluronidase 0.05% for 15 minutes, trypsin 0.25% for 30 minutes and collagenase 0.2% for 90 minutes. Finally, the HAM's F12 medium is
20 added for 60 minutes. The suspension of chondrocytes is cultured in dishes 6 cm in diameter (150,000 cells per dish) in the presence of HAM's F12 medium containing 10% fetal calf serum. Under these conditions, the chondrocytes have a differentiated phenotype as the production of type II collagen demonstrates.

- When these cells reach preconfluence (6 to 7 days), they are transfected with
25 the various constructs using the calcium phosphate method.

2- Measurement of the CAT and β -galactosidase activities.

The CAT activity is measured according to the double liquid phase technique according to Desbois et al. (1992). The β -galactosidase activity is measured with the aim of normalizing the variations in the transfection efficiency.

3- Preparation of the nuclear proteins and retardation experiments in electrophoresis.

The Cos-1 cells (transformed monkey kidney fibroblasts) are transfected according to the abovementioned calcium phosphate technique using 20 μ g of vectors for expression of PPAR γ , PPAR α or RXR. Non-transfected Cos-1 cells will serve as control. 48 hours after the transfection, the cells are incubated with a buffer containing KCl (0.45 M), Tris-HCl (20 mM, pH=7.5) and 10% glycerol. They are detached by scraping and then disrupted by cooling to -80°C and thawing at 37°C . These cells are then centrifuged at 15,000 rpm, for 10 minutes at 4°C . The supernatant is then frozen at -80°C and used for the gel retardation experiments.

4- Gel retardation experiments.

The sequences DR1, (DR1) $_2$ 17, (DR1) $_2$ 21, and (DR1) $_2$ 31 are radiolabeled by means of the Klenow fragment of DNA polymerase using [α - ^{32}P]dCTP. One hundred thousand cpm (0.4 ng) of the radiolabeled probes are incubated, for 30 minutes at 4°C , with increasing quantities of Cos-1 extracts enriched with PPAR and RXR (as indicated in the legend to the figures) and with a specific binding buffer (KCl 100 mM, Hepes 40 mM, pH 7.5, NP40 0.1%, ZnCl_2 0.5 mM, PEG 8%). The mixture is then deposited on an acrylamide/bisacrylamide gel (5%, TBE 0.25 X) and run for 2 hours 30 min at 200 volts. The gel is then dried and autoradiographed.

5- Molecular biology

5 The methods commercially used in molecular biology such as preparative extractions of plasmid DNA, centrifugation of plasmid DNA in a cesium chloride gradient, electrophoresis on agarose gels, purification of DNA fragments by electroelution, precipitation of plasmid DNA in saline medium with ethanol or isopropanol, transformation in Escherichia coli are well known to persons skilled in the art and are widely described in the literature (Sambrook et al., "Molecular Cloning, a Laboratory Manual", Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1989).

10 The plasmids are of commercial origin.

The enzymatic amplification of the DNA fragments by the PCR technique (Polymerase Chain Reaction) may be carried out using a DNA thermal cyclerTM (Perkin Elmer Cetus) according to the manufacturer's recommendations.

15 The electroporation of plasmid DNA into Escherichia coli cells may be carried out with the aid of an electroporator (Bio-Rad) according to the manufacturer's recommendations.

20 The verification of the nucleotide sequences may be carried out by the method developed by Sanger et al. (*Proc. Natl. Acad. Sci. USA*, 74 (1977) 5463-5467) using the kit distributed by Applied Biosystems according to the manufacturer's recommendations.

EXAMPLES

25 In the experimental section which follows, there have been described the construction and use of several hybrid promoters of the invention, comprising an element inducible in particular by IL1b and an element inducible by icosanoids, fatty acids and nonsteroidal anti-inflammatory agents (NSAID).

The first module comprises in particular the region [-247; +20] (relative to the site of initiation of transcription of sPLA2). This region contains the TATA box and the proximal element (-114/-85) which is essential for the transcriptional activity and which binds the ubiquitaire factor Sp1. It also comprises the site C/EBP (-198/-190) essential for stimulation by IL-1 β and the element [-247/-210] which enhances the activity of the C/EBP factors. Furthermore, this module contains a glucocorticoid response element capable of relaying induction by these hormones.

The addition of the second module is intended to enhance the induction during the inflammatory process by the mediators derived from the fatty acids produced by the inflammation (prostaglandins, leukotrienes) or by exogenous compounds (fibrate, nonsteroidal anti-inflammatory drugs-NSAID). This module comprises a PPARs response element (PPRE) situated upstream of the first module.

EXAMPLE 1: Construction of PPAR response elements

This example describes the construction and functional analysis of several PPAR response elements which can be used in the hybrid promoters of the invention.

1- Production of a PPARs response unit of high affinity

In order to obtain optimum induction by the PPAR ligands, several arrangements of PPRE multimers were designed and synthesized. The sequence of these regions is given below.

The element DR1 is the consensus site for binding of the heterodimer PPAR/RXR. The elements (DR1)₂ 17, (DR1)₂ 21 and (DR1)₂ 31 consist of two DR1s separated center to center by 17, 21 and 31 base pairs respectively.

DR1 (SEQ ID NO: 1)

CAAACT **AGGTCA** A **AGGTCA**

(DR1)₂ 17 (SEQ ID NO:2)

CAAACT **AGGTCA** A **AGGTCA** AACT **AGGTCA** A **AGGTCA**

(DR1)₂ 21 (SEQ ID NO: 3)

CAAACT **AGGTCA** A **AGGTCA** t CAAACT **AGGTCA** A **AGGTCA**

(DR1)₂ 31 (SEQ ID NO: 4)

CAAACT **AGGTCA** A **AGGTCA** tgtctttaggcc CAAACT **AGGTCA** A **AGGTCA**

2- Binding experiments

2-1 binding of PPAR_γ

In order to verify the functional nature and the efficiency of the above PPRE regions, gel retardation experiments were carried out consisting in studying the binding between one of the four sequences above, radiolabeled beforehand, in the presence of increasing quantities (5 and 10 μ l) of PPAR_γ and RXR proteins prepared in Cos-1 cells (PPAR_γ/RXR ratio=3:1). The results obtained are presented in Figure 2.

These results show that the sequences constructed bind the heterodimer PPAR/RXR. The sequence DR1 binds a dimeric complex whereas the sequences (DR1)₂ 17, (DR1)₂ 21 and (DR1)₂ 31 bind the dimeric complex but also a retarded

complex of high molecular weight which is thought to correspond to a double heterodimer PPAR/RXR. Moreover, the quantities of the dimeric and high-molecular weight complexes are variable from one sequence to another. This suggests that these three sequences bind the RXR/PPAR γ complexes with a different cooperativity.

To know if the complexes observed come from the production of PPAR γ and from RXR in the Cos-1 cells, or alternatively from proteins present in these cells, these different sequences were incubated with extracts of nontransfected Cos-1 (10 μ l). The intensity of the complexes formed with these extracts is very weak, which is in favor of the production of PPAR γ and RXR proteins following the transfection of the cells (Fig. 2).

With the aim of studying the cooperativity of the binding of PPAR/RXR to these different sequences, gel retardation experiments placing increasing quantities of Cos-1 extracts enriched with PPAR and RXR (1 μ l to 10 μ l), followed by quantification of the radioactivity obtained, were carried out. The formation of the first PPAR/RXR dimer is characterized by the dissociation constant KdII, that of the double dimer, or tetramer, is characterized by the dissociation constant KdIV. The KdII/KdIV ratio (R) is the reflection of the cooperativity of the binding if it is greater than 1.

The various dimeric and tetrameric complexes and the free probe were quantified by means of a Phosphorimager and represented in Figure 3. The cooperativity ratios were calculated by means of the abovementioned formula. The sequences (DR1)₂ 17 and (DR1)₂ 31 bind as many dimeric and tetrameric forms. The cooperativity ratios are 9 and 12.6 respectively. However, the sequence (DR1)₂ 21 exhibits a different binding profile. The dimeric form forms only 20% of the total

binding, whereas the tetrameric form forms 80% of the total binding. The cooperativity ratio is 34 +/- 11.

These results as a whole show that the three sequences used cooperatively bind PPAR and RXR. The sequence which exhibits the best cooperativity of binding is (DR1)₂ 21.

2-2 binding of PPAR α

Similar experiments were carried out with PPAR α . The vector for expressing the α isoform of PPAR was transfected into Cos-1 cells. Total extracts of the latter were prepared as described in the section Materials and Methods. Gel retardation experiments using the four radiolabeled sequences and increasing quantities of Cos-1 extracts enriched with PPAR α and RXR (rapport3/1) were carried out. The bands corresponding to the dimeric (D) or tetrameric (T) forms or the free probe (P) were cut out of the gel and then quantified by counting on a beta scintillation counter (Figure 4). The thermodynamic model described above was applied with the aim of calculating the ratios of cooperativity of binding of the PPAR α /RXR heterodimer to the different sequences studied.

By contrast to what was observed with PPAR γ , the cooperativity ratios obtained with the sequences (DR1)₂ 21 and (DR1)₂ 31 are equivalent (R=11.2 and 11.6 respectively). The cooperativity ratio obtained in the case of the element (DR1)₂ 17 is less than the preceding two (R=7). In this case also, the three synthesized sequences cooperatively bind PPAR α , the sequences (DR)₂ 21 and (DR1)₂ 31 exhibiting the best binding cooperativity.

EXAMPLE 2: Construction and functional analysis of a vector comprising a hybrid promoter.

This example describes construction of plasmids comprising a hybrid promoter of the invention, and demonstrates its induction properties by various inflammation mediators.

1) Construction of the plasmids and functional tests

The fragment -247/+20 of the promoter of sPLA2 IIA (SEQ ID NO : 5) was subcloned upstream of the CAT reporter vector for the plasmid PUC-SH-CAT. Upstream of this promoter, the different PPRE sequences cited above (SEQ ID NO : 1-4) were inserted.

The rabbit chondrocytes were then transfected with the CAT constructs containing the different arrangements of PPRE (DR1) placed upstream of the minimal promoter of sPLA2-IIA. 15 deoxy Δ^{12-14} PGJ2 (10 μ M) (activator of PPAR γ) or Wy-14643 (200 μ M) (activator of PPAR α) were then added to the culture medium for 24 hours. The results obtained are presented in Figure 5.

The results show that the minimal promoter (-247/+20) of sPLA2-IIA, which does not contain known PPRE sites, is not activated by PGJ2 or by Wy 14643. However, the different arrangements of DR1 are activated at different levels by the inducers of PPAR γ or of PPAR α . It is possible to classify them in the following increasing order: DR1 < (DR1)_{2 17} = (DR1)_{2 31} < (DR1)_{2 21} in the case of PPAR γ . On comparing the binding results with the functional results, it is observed that the

sequence (DR1)_{2 21}, which has the best binding cooperativity, exhibits the best functional synergy.

In the case of PPAR α , the order of inducibility by Wy-14643 is different from that established in the case of PGJ2. DR1 < (DR1)_{2 21} < (DR1)_{2 17} < (DR1)_{2 31}. The gel retardation experiments are not in perfect agreement with the transient transfection experiments. The binding experiments showed that the sequences (DR1)_{2 21} and (DR1)_{2 31} bind PPAR α with the same cooperativity (R=11) whereas the sequence (DR1)_{2 17} exhibits a lower binding cooperativity (R=7).

2- Combined effects of IL-1, of glucocorticoids and of inducers of PPAR α (fibrates) and PPAR γ (15 dioxo Δ 12-14 PGJ2) on the synthetic promoter:

Rabbit chondrocytes were then transfected, either with the minimal promoter of sPLA2-IIA, or with the sequence (DR1)_{2 21} placed upstream of the latter (hybrid promoter). PGJ2 (10 μ M) and/or IL-1 (10 ng/ml) are added to the culture medium.

The results obtained are presented in Figure 6. These results show that IL-1 induces the promoter of sPLA2-IIA 6 fold. The prostaglandin J2 or Wy-14643 have no effect on this promoter (results not shown). In the case of the construct (DR)_{2 21}-(-247/+20) sPLA2-CAT, IL-1 and PGJ2 induce this promoter 4 fold. The two combined products have a synergistic effect (13-fold induction relative to the basal activity) (Fig. 6).

This same type of experiment was carried out with Wy-14643. This compound is an inducer specific for PPAR α . Wy-14643 (200 μ M) activates the transcription of the construct (DR)_{2 21}-(-247/+20) sPLA2-CAT about 3 to 4 fold. The effect of the combined IL-1 and Wy-14643 is synergistic (20-fold induction).

The effect of dexamethasone was moreover tested on the synthetic promoter (-247/+20)-(DR1)₂ 21-CAT, either alone or combined with Wy-16463, PGJ2, or IL-1. Dexamethasone activates the transcription of this promoter by about 6 fold. These results show that the circulating glucocorticoids such as cortisol, or administered during a treatment, can activate this hybrid promoter.

The effect of some nonsteroidal anti-inflammatory drugs such as indomethacin or ibuprofen was also tested on the synthetic promoters. The results obtained show that indomethacin (10^{-4} M) activates by about fivefold the construct (DR1)₂ 21-(-247/+20)-CAT whereas ibuprofen (10^{-4} M) activates this promoter 17 fold.

Finally, the levels of transcriptional activities reached by a viral promoter, such as the RSV (Rous Sarcoma Virus) promoter, and that of the synthetic promoter were compared. The basal transcriptional activity of the synthetic promoter is low, it is less than 5% of the activity of the RSV viral promoter. After induction with the different effectors, the synthetic promoter exhibits a level of transcriptional activity which reaches 40% of that of RSV (results not shown). The promoter constructed according to the present invention is therefore characterized by a high induced transcriptional activity comparable to the viral vectors currently used in gene therapy, and a low basal activity.

Conclusions

Hybrid synthetic promoters inducible by the components of inflammation, that is to say the pro-inflammatory cytokines such as IL-1 β , as well as the pro-inflammatory fatty acids, such as the prostaglandins or the leukotrienes were developed. Furthermore, the circulating or locally administered glucocorticoids as well as the nonsteroidal anti-inflammatory drugs, such as ibuprofen or indomethacin,

are capable of activating the transcription of these promoters. These promoters therefore constitute an effective means of modulating the activity of genes according to whether an anti-inflammatory treatment is administered or not. These characteristics make it possible to envisage a controllable (on/off) effect on the synthesis of the medicament gene. It is important to note that this synthetic promoter has a very low basal activity, whereas it has a transcriptional activity induced by the endogenous or exogenous stimuli comparable to the viral vectors in efficiency.

5

5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
22

BIBLIOGRAPHIC REFERENCES

- Amin AR, Attur M, Patel RN, Thakker GD, Marshall PJ, Rediske J, Stuchin SA, Patel IR, Abramson SB (1997) Superinduction of cyclooxygenase-2 activity in human osteoarthritis-affected cartilage. Influence of nitric oxide. *J Clin Invest* 99 (6): 1231-1237.
- 5 Berenbaum F, Thomas G, Poiraudau S, Bereziat G, Corvol MT, Masliah J (1994) Insulin-like growth factors counteract the effect of interleukin 1 beta on type II phospholipase A₂ expression and arachidonic acid release by rabbit articular chondrocytes. *FEBS Lett* 340 (1-2): 51-55.
- 10 Bingham C. O. 3rd, Murakami M., Fujishima H., Hunt J. E., Austen K. F., Arm J. P. (1996) A heparin-sensitive phospholipase A₂ and prosaglandin endoperoxide synthase-2 are functionally linked in the delayed phase of prostaglandin D₂ generation in mouse bone marrow-derived mast cells. *J. Biol. Chem.* 271: 25936-25944.
- 15 Camper L, Heinegard D, Lundgren-Akerlund E (1997) Integrin alpha2beta1 is a receptor for the cartilage matrix protein chondroadherin. *J Cell Biol* ;138(5):1159-6
- Chevalier, X. *Physiopathologie de l'arthrose [Physiopathology of osteoarthritis]*. (1998) La Presse Médicale. 27(2). 75-92.
- Clark JD, Lin LL, Kriz RW, Ramesha CS, Sultzman LA, Lin AY, Milona N, Knopf J (1991) A novel arachidonic acid-selective cytosolic PLA₂ contains a Ca²⁺ dependent translocation domain with homology to PKC and GAP. *Cell* 65: 1043-1051.
- 20 Dean, J, Martel-Pelletier, J, Pelletier, JP, et al. Evidence for metalloprotease and metalloprotease inhibitors imbalance in human osteoarthritic cartilage. *Arthritis Rheum.* 1990, 33, 1466-76
- Desbois C, Massé T, Madjar JJ. (1992) Optimization of the CAT assay procedure by determining the initial rate of the enzymatic reaction. *Trends Genet*; 8 : 300-30
- 25 Devchand, RP, Keller, H, Peters, JM, Vazquez, M, Gonzalez, FJ, Wahli, W. The PPAR ?-leukotriene B₄ pathway to inflammation control. (1996) *Nature*, 384, 39-43.

Di Battista JA, Dore S, Martel-Pelletier JP (1996) Prostaglandin E2 stimulates incorporation of proline into collagenase digestible proteins in human articular chondrocytes: identification of an effector autocrine loop involving insulin-like growth factor I. *Mol Cell Endocrinol* 123: 27-35.

- 5 Geng Y, Blanco FJ, Cornelisson M, Lotz M (1995) Regulation of cyclooxygenase-2 expression in normal human articular chondrocytes. *J Immunol* 155 (2): 796-801

Guingamp C, Gegout-Pottie, Philippe L, Terlain B, Netter P, Gillet P. (1997) Mono-iodoacetate-induced experimental osteoarthritis. *Arthritis Rheumatism*. 40:1670-1679.

Hornebeck W, Lafuma, C. Les métalloprotéinases matricielles [Matrix metalloproteinases].

- 10 (1990) *C R Soc biol*. 185, 1466-76

Hulkower KI, Hope WC, Chen T, Anderson CM, Coffey JW, Morgan DM (1992) Interleukin-1? stimulates cytosolic phospholipase A₂ in rheumatoid synovial fibroblasts. *Biochem. Biophys. Res. Commun*. 184: 712-718.

Jacques, C., Bereziat, G., Humbert, L., Olivier, J. L., Corvol, M. T., Masliah, J., and

- 15 Berenbaum, F. (1997). Posttranscriptional effect of insulin-like growth factor-I on interleukin-1beta-induced type II-secreted phospholipase A₂ gene expression in rabbit articular chondrocytes. *J Clin Invest* 99, 1864-72.

Jpenberg AI, Jeannin E, Wahli W, Desvergne B. (1997) Polarity and specific sequence requirements of peroxisome proliferator-activated receptor (PPAR)/retinoid X receptor
20 heterodimer binding to DNA.

Kim Y, Fischer SM (1998) Transcriptional regulation of cyclooxygenase-2 in mouse skin carcinoma cells. Regulatory role of CCAAT/enhancer-binding proteins in the differential expression of cyclooxygenase-2 in normal and neoplastic tissues. *J Biol Chem* 16;273(42):27686-9

- 25 Klapisz E, Ziari M, Wendum L, Koumanov K, Brachet C, Ducos, Olivier JL, Béréziat G, Trugnan G, Masliah J (1999) N- and C-terminal plasma membrane anchoring modulate differently agonist-induced activation of cytosolic phospholipase A₂ (publication en cours).

Kliwer, SA, Lenhard, JM, Wilson, TM, Patel, I, Morris, DC, Lehmann, JM. A prostaglandine J2 metabolite binds peroxisome proliferator-activated receptor ? and promotes adipocyte differentiation. (1995) *Cell*. 83, 813-819

Knott I, Dieu M, Burton M, Houbion A, Remacle J, Raes M (1994) Induction of cyclooxygenase by interleukin 1: comparative study between human synovial cells and chondrocytes. *J Rheumatol* 21 (3): 462-466.

Kreiss P, Scherman D (1999) Optimisation des plasmides et des vecteurs synthétiques pour la thérapie génique [Optimization of the synthetic vectors and plasmids for gene therapy]. *Médecine/Sciences*, 15 : 669-676

Krey, G, Braissant, O, L'Horset, F, Kalkhoven, E, Perroud, M, Parker, MG, Wahli, W. Fatty acids, eicosanoids, and hypolipidemic agents identified as ligands of peroxisome proliferator-activated receptors by coactivator-dependent receptor ligand assay. (1997) *Mol. Endocrinol.* 11, 779-791.

Kuwata H., Nakatani Y., Murakami M. & Kudo I. (1998) Cytosolic phospholipase A₂ is required for cytokine-induced expression of type IIA secretory phospholipase A₂ that mediates optimal cyclooxygenase-2-dependent delayed prostaglandin E₂ generation in rat 3Y1 fibroblasts. *J. Biol. Chem.* 273: 1733-40.

Leevers SJ, Paterson HF, Marshall CJ (1994) Requirement for Ras in Raf activation is overcome by targeting Raf to the plasma membrane. *Nature* 369, 411-414.

Lefebvre, V & de Crombrughe, B. Toward understanding SOX9 function in chondrocyte differentiation. *Matrix Biol.* 16, 529-540

Lefebvre, V, Huang, W, Harley, VR, Goodfellow, PN & de Crombrughe, B. SOX9 is a potent activator of the chondrocyte-specific enhancer of the pro alpha1(II) collagen gene. (1997) *Mol. Cell. Biol.* 17, 2336-2346

Lehmann JM, Lenhard JM, Oliver BB, Gordan MR, Kliwer SA. (1997). Peroxisome Proliferator-activated receptor α and are activated by indomethacin and other non-steroidal anti-inflammatory drugs. *J. Biol. Chem.* 272:3406-3410.

- Lin M. K., Farewell V., Vadas P., Bookman A. A., Keystone E. C., Pruzanski W. (1996) *Secretory phospholipase A2 as an index of disease in rheumatoid arthritis. Prospective double blind study of 212 patients. J. Rheumatol. 23: 112-1166.*
- Makihira S, Yan W, Ohno S, Kawamoto T, Fujimoto K, Okimura A, Yoshida E, Noshiro M, Hamada T, Kato Y. Enhancement of cell adhesion and spreading by a cartilage-specific noncollagenous protein, cartilage matrix protein (CMP/Matrilin-1), via integrin $\alpha 1 \beta 1$. (1999) *J Biol Chem* , 274(16):11417-2
- Mukhopadhyay, K, Lefebvre V, Zhou, G, Garofalo, S, Kimura, JH and de Crombrügge, B. Use of a new rat chondrosarcoma cell line to delineate a 119-base pair chondrocyte-specific enhancer element and to define active promoter segments in the mouse pro- $\alpha 1$ (II) collagen gene (1995) *J. Biol. Chem.* 270, 27711-19.
- Nevalainen TJ, Marki F, Kortesuo PT, Grutter MG, Di Marco S, Schmitz A (1993) Synovial type (group II) phospholipase A_2 in cartilage. *J Rheumatol* 20 (2): 325-330.
- Pruzanski W., Albin-Cook K., Laxer R. M., MacMillan J., Stefanski E., Vadas P., and Silverman E. D. (1994) Phospholipase A_2 in juvenile rheumatoid arthritis: Correlation to disease type and activity. *J. Rheumatol.* 21: 1951-1954.
- Schoonjans, K, Staels, B, Auwrex, J. The peroxisome proliferator activated receptors (PPARs) and their effects on lipid metabolism and adipocyte differentiation. (1996) *Biochem. Biophys. Acta.* 1302, 93-109.
- Seilhamer JJ, Pruzanski W, Vadas P., Plant S, Miller JA, Kloss J, Johnson LK (1989) Cloning and recombinant expression of phospholipase A_2 present in rheumatoid arthritic synovial fluid. *J. Biol. Chem.* 264: 5335-5338.
- Stokoe D, Macdonald SG, Cadwallader K, Symons M, Hancock JF (1994) Activation of Raf as a result of recruitment to the plasma membrane. *Science* 264, 1463-1467.
- Thomas B, Humbert L, Crofford L, Biu X, Berenbaum F, Olivier JL. Critical role of C/EBPd and C/EBPb factors in the stimulation of cyclooxygenase-2 gene transcription by interleukine-1b in primary culture chondrocyte (publication pending).

Zhou, G, Garofalo, S, Mukhopadhyay, K, Lefebvre, V, Smith, CN, Eberspaecher, H and de Crombrugge, B. A 182 bp fragment of the mouse alpha 1 (II) collagen gene is sufficient to direct chondrocyte expression in transgenic mice (1995) *J. Cell. Sci.* 108, 3677-84.

Zlatkine P, Mehul B, Magee AI (1997) Retargeting of cytosolic proteins to the plasma membrane by the Lck protein tyrosine kinase dual acylation motif. *J. Cell. Science* 110, 673-679.